



# Occurrence of *Vernonia anamallica* (Asteraceae) in Agasthyamalai Biosphere Reserve, southern Western Ghats, India

R. Jagadeesan<sup>1</sup>, A. Gangaprasad<sup>1\*</sup> and S.P. Mathew<sup>2</sup>

<sup>1</sup>Department of Botany, University of Kerala, Kariavattom P.O., Thiruvananthapuram – 695562, Kerala, India.

<sup>2</sup>Jawaharlal Nehru Tropical Botanic Garden and Research Institute (JNTBGRI), Palode, Karimankode P.O., Thiruvananthapuram – 695562, Kerala, India.

\*E-mail: agangaprasad@yahoo.com

## Abstract

*Vernonia anamallica* Bedd. ex Gamble was described based on the collection from the Anamalai hills, southern Western Ghats. This species was remained unknown until it was rediscovered after 92 years from the Rajamala hills and subsequently known by two more collections from Pettimudi and Konalar. A detailed description along with photographs is provided here for the first time based on our recent collection from the Agasthyamalai Biosphere Reserve (ABR), which also forms an extended distribution of this species.

**Keywords:** Agasthyamalai Biosphere Reserve, Endemic, Southern Western Ghats, *Vernonia anamallica*

## Introduction

The genus *Vernonia* was established by a German naturalist, Johann Christian Daniel von Schreber in 1791. It is represented by c. 685 taxa (www.theplantlist.org) and distributed both in Old and New Worlds with centres of diversity in East Africa and Brazil (Keeley *et al.*, 2007, 2009). Hooker (1881) enumerated a total of 45 species and 3 varieties from the Indian subcontinent. Subsequently, Uniyal (1995) and Karthikeyan *et al.* (2009) have reported 52 species and 10 varieties and 55 species, 1 subspecies and 7 varieties, respectively from India. Nayar *et al.* (2014) reported 17 taxa, including 12 endemics from the Western Ghats.

Gamble (1920) described *Vernonia anamallica* based on the collection of R.H. Beddome made during 1873 from the Anamalai hills in southern Western Ghats. The Beddome's collection was originally with two specimens on same sheet; however, Gamble had removed one of them and annotated on the type sheet as "I have taken the liberty of keeping the upper specimen for Kew – please excuse" and incorporated this specimen (K000814740) at K. This species was remained unknown until it was rediscovered after more than 92 years from Rajamala hills in 1965 and Pettimudi in 1970 by Shetty & Vivekanathan (1972) without any illustrations or photographs. It was also collected from Konalar in the Anamalais of Tamil Nadu during the early and late 1980s

by Chandrabose. During the recent exploration tour to Agasthyamalai Biosphere Reserve, two individuals of this species were found, an evidence of its extended distribution from the earlier known localities. Therefore, a brief description of the species along with colour photographs is provided here for the first time to facilitate its precise identification.

***Vernonia anamallica*** Bedd. ex Gamble, Bull. Misc. Inform. Kew 1920: 339. 1920 & Fl. Madras: 675. 1921; B.V. Shetty & Vivek., Bull. Bot. Surv. India 14: 19. 1972; Uniyal in Hajra *et al.*, Fl. India 13: 353, f. 110. 1995; Karthik. *et al.*, Fl. Pl. India 1: 291. 2009. Type: INDIA, **Tamil Nadu**, Coimbatore district, Anamalai hills, in higher ranges, 1873, R.H. Beddome s.n. (K000814740, image!). **Fig. 1**

Shrub, to 2 m high. Stems angled with prominent nodal scars, woody; branches appressed-hairy when young, glabrous on maturity. Leaves simple, opposite, elliptic, elliptic-lanceolate or elliptic-ovate, 3.5–10.5 × 1.5–5 cm, shortly attenuate, rarely oblique at base, serrate to crenate at margins, acuminate, rarely mucronate at apex, pale greenish above, golden pubescent beneath; lateral veins 5–8 pairs, convergent towards apex, conspicuous beneath; petioles 0.5–1 cm long. Synflorescence corymbose, c. 4 cm long; peduncles 2–3 cm long. Heads c. 1 cm long; phyllaries ovate, c. 3 × 1.5 mm, mucronate at apex, tomentose. Florets 5–7 mm long. Corolla tubular, white, 5-lobed; lobes



Fig. 1. *Vernonia anomallica* Bedd. ex Gamble: a. Flowering Twig; b. Head; c. Floret; d. Corolla split-open; e. Anthers; f. Pistil; g. Phyllary; h. Achene; i. Type specimen (R.H. Beddome s.n., K000814740).

lanceolate, 2–4 mm long. Stamens 5; filaments short; anthers c. 2.5 mm long, purple; style linear, white; stigma bifurcated, arching down. Achenes 5-angled, glabrous; pappus 2-seriate.

*Flowering & Fruiting*: November–March.

*Habitat*: Moist rocky localities near shola forest, 1580–2075 m. Only two individuals were found growing along with *Phyllanthus macraei* Müll.Arg., *Ixora undulata* Roxb., *Aenhenrya rotundifolia* (Blatt.) C.S. Kumar & F.N. Rasm., *Exacum travancoricum* Bedd. and *Crotalaria scabra* Gamble.

*Distribution*: India (Kerala and Tamil Nadu). **Endemic.**

*Specimens examined*: INDIA, **Kerala**, Idukki district, Rajamallay, 2075 m, 21.11.1965, B.V. Shetty 26598 (MH); Pettimudi, 1675 m, 8.2.1970, B.V. Shetty 33450 (MH); Thiruvananthapuram district, Agasthyamalai Biosphere Reserve (Druiry hills), 1580 m, 23.12.2015, R. Jagadeesan 8020 (KUBH, TBGT). **Tamil Nadu**, Coimbatore district, Konalar, 15.2.1980, 1850 m, M. Chandrabose 65816 (MH).

## Acknowledgements

The authors are thankful to the Head, Department of Botany, University of Kerala and the Director, JNTBGRI, Thiruvananthapuram, for facilities.

## Literature Cited

- Gamble, J.S. 1920.** Decades Kewensis – LIV. Plantarum Novarum in herbario horti regii conservatarum. Decades C–CI. *Bull. Misc. Inform. Kew* **1920**: 339. 1920.
- Hooker, J.D. 1881.** Compositae. *The flora of British India*. Vol. 3. L. Reeve & Co. Ltd., London. pp. 219–419.

**Karthikeyan, S., Sanjappa, M. & S. Moorthy 2009.** *Flowering plants of India – Dicotyledons*. Vol. 1 (Acanthaceae–Avicenniaceae). Botanical Survey of India, Kolkata. pp. 184–299.

**Keeley, S.C. & H. Robinson 2009.** Vernonieae. In: Funk, V.A., Susanna, A., Steussy, T.F. & R.J. Bayer (eds.), *Systematics, Evolution, and Biogeography of Compositae*. International Association for Plant Taxonomy, University of Vienna, Vienna. pp. 439–461.

**Keeley, S.C., Forsman, Z.H. & R. Chan 2007.** A phylogeny of the “evil tribe” (Vernonieae: Compositae) reveals Old/New World long distance dispersal: Support from separate and combined congruent datasets (trnL–F, ndhF, ITS). *Molec. Phylogen. Evol.* **44**: 89–103.

**Nayar, T.S., Rasiya Beegam, A., Mohanan, N. & G. Rajkumar 2006.** *Flowering plants of Kerala: A handbook*. Tropical Botanic Garden and Research Institute, Thiruvananthapuram.

**Shetty, B.V. & K. Vivekananthan 1972.** New and little known taxa from Anaimudi and surrounding regions, Devicolam, Kerala – IV: Notes on some rare species. *Bull. Bot. Surv. India* **14**: 19–23.

**Uniyal, B.P. 1995.** Vernonieae. In: Hajra, P.K., Rao, R.R., Singh, D.K. & B.P. Uniyal (eds.), *Flora of India*. Vol. 13. Asteraceae (Inuleae–Vernonieae). Botanical Survey of India, Calcutta. pp. 330–394.

---

Received: 30.5.2016

Revised and Accepted: 29.11.2016